arXiv:1210.1942v1 [math.NT] 6 Oct 2012

IDENTITIES FOR THE RAMANUJAN ZETA FUNCTION

MATHEW ROGERS

Abstract. We derive formulas for special values of the Ramanujan zeta function outside of the critical strip. In particular, we give explicit integrals for $L(\Delta,12)$ and $L(\Delta,13)$ which involve powers of hypergeometric functions. Our method also shows that $L(\Delta,k)$ is a period in the sense of Kontsevich and Zagier when $k\geq 12$.

1. Introduction

Ramanujan introduced his zeta function in 1916 [6]. Following Ramanujan, let

$$\Delta(z):=q\prod_{n=1}^{\infty}(1-q^n)^{24}=\sum_{n=1}^{\infty}\tau(n)q^n,$$

where $q=e^{2\pi iz}$. Since $\tau(n)$ is multiplicative, Ramanujan had the natural idea to look at the following Dirichlet series:

$$L(\Delta,s):=\sum_{n=1}^{\infty}\frac{\tau(n)}{n^s}.$$

Ramanujan observed that $L(\Delta,s)$ has an Euler product, and it was subsequently proved that the function analytically continues to the complex plane [10, pg. 242]. The function satisfies a relation of the form $L(\Delta,12-s)=\gamma(s)L(\Delta,s)$, where $\gamma(s)$ involves Gamma functions. The main goal of this paper is to prove formulas for $L(\Delta,k)$ when $k\geq 12$ is a positive integer lying outside of the critical strip. We use recent ideas of Rogers and Zudilin [8], [9], and Zudilin [11], to reduce these L-values to elementary integrals. For example, in Theorem 1 we prove

$$\begin{aligned}L(\Delta,12)=-\frac{128\pi^{11}}{8241\cdot 11!}\int_0^1&[F(\alpha)F(1-\alpha)]^5\\&\times\left(\frac{2+251\alpha+876\alpha^2+251\alpha^3+2\alpha^4}{1-\alpha}\right)\log\alpha\ \mathrm{d}\alpha,\end{aligned}\tag{1}$$

where $F(\alpha)$ is the usual hypergeometric series:

$$F(\alpha):={}_2F_1\left({\tfrac12,\tfrac12 \atop 1};\alpha\right)=\sum_{n=0}^{\infty}\binom{2n}{n}^2\left(\frac{\alpha}{16}\right)^n.\tag{2}$$

Date: October 4, 2012.

2010 *Mathematics Subject Classification.* Primary 33C20; Secondary 11F11, 11F03, 11Y60, 33C75, 33E05.

Key words and phrases. Ramanujan zeta function, L-values.

We prove a double integral for $L(\Delta,13)$ in Theorem 2, and we point out that it is possible to obtain similar double integrals for $L(\Delta,k)$ when $k\geq 13$.

2. Critical values of $L(\Delta,s)$

We begin with a brief review of known formulas for critical values of $L(\Delta,s)$. We say that $L(\Delta,k)$ is a critical L-value if $1\leq k\leq 11$. Kontsevich and Zagier summarized the properties of critical L-values in [4]. It is relatively simple to relate these values to integrals of algebraic functions. The standard Mellin transform gives

$$L(\Delta,k)=\frac{(2\pi)^k}{(k-1)!}\int_0^\infty u^{k-1}\Delta(iu)\mathrm{d}u \tag{3}$$

whenever $k\geq 1$. The usual method for obtaining an elementary integral, is to set

$$u=\frac{F(1-\alpha)}{2F(\alpha)}, \tag{4}$$

where $F(\alpha)$ is defined in (2). Notice that $\tau=iu$ is the period ratio of the elliptic curve $y^2=(1-x^2)(1-\alpha x^2)$. Now appeal to the classical formulas [2, pg. 124], [2, pg. 120]:

$$\Delta(iu)=\frac{1}{16}\alpha(1-\alpha)^4\left[F(\alpha)\right]^{12}, \tag{5}$$

$$\frac{\mathrm{d}u}{\mathrm{d}\alpha}=\frac{-1}{2\pi\alpha(1-\alpha)\left[F(\alpha)\right]^2}, \tag{6}$$

and notice that $\alpha\in(1,0)$ when $u\in(0,\infty)$. Equation (3) reduces to

$$L(\Delta,k)=\frac{\pi^{k-1}}{16(k-1)!}\int_0^1(1-\alpha)^3\left[F(\alpha)\right]^{11-k}\left[F(1-\alpha)\right]^{k-1}\mathrm{d}\alpha. \tag{7}$$

This formula illustrates the crucial point that $L(\Delta,k)$ belongs to the *ring of periods* when $1\leq k\leq 11$. In other words, $L(\Delta,k)$ can be written as a multiple integral of purely algebraic functions. We can make this explicit by substituting a classical elliptical integral for every hypergeometric series:

$$F(\alpha)=\frac{2}{\pi}\int_0^1\frac{\mathrm{d}u}{\sqrt{(1-u^2)(1-\alpha u^2)}}. \tag{8}$$

Kontsevich and Zagier illustrated this point with an equivalent version of (7), albeit in a slightly different notation [4, pg. 24].

3. A formula for $L(\Delta,12)$

It is *extremely non-obvious* that $L(\Delta,k)$ is a period if $k\geq 12$. Equation (7) becomes an integral over non-trivial ratios of hypergeometric functions, and it is no longer clear how to simplify the identity. While Beilinson's theorem implies that $L(\Delta,k)$ is a period outside of the critical strip [1], it is quite challenging to use his result to extract explicit formulas. The goal of this section is to prove a formula for $L(\Delta,12)$ using the method that Rogers and Zudilin introduced in [8] and [9]. In Section 4 we study $L(\Delta,k)$ when $k\geq 13$.

Zudilin has provided a nice summary of our method in [11]. The crucial step is to decompose $\Delta(z)$ into a linear combination of products of two Eisenstein series. The most famous decomposition of $\Delta(z)$ is due to Ramanujan:

$$1728\Delta(z) = E_4(z)^3 - E_6(z)^2, \tag{9}$$

where

$$E_4(z) := 1 + 240\sum_{n=1}^{\infty}\frac{n^3q^n}{1-q^n}, \qquad E_6(z) := 1 - 504\sum_{n=1}^{\infty}\frac{n^5q^n}{1-q^n},$$

and $q = e^{2\pi iz}$. Equation (9) is not helpful in this context, because it involves a cubic power $E_4(z)$. In Lemma 1, we decompose a linear combination of $\Delta(z)$'s into the desired form.

Lemma 1. *We have*

$$\begin{aligned}\Delta(z) + 24\Delta(2z) + 2^{11}\Delta(4z) = \frac{8}{504^2}&\left[E_6(2z) - 64E_6(4z)\right] \\ &\times\left[E_6(z) - 33E_6(2z) + 32E_6(4z)\right].\end{aligned} \tag{10}$$

Proof. We used a computer to check that the first 100 Fourier coefficients of both sides agree. Since both functions are modular forms on $\Gamma_0(4)$, we can use Rankin's theorems to conclude that they are equal [7]. For a nice discussion of this method see [2, pg. 334]. ☐

Now apply an involution to the first term on the right-hand side of (10). The equation becomes

$$\begin{aligned}\Delta(z) + 24\Delta(2z) + 2^{11}\Delta(4z) = \frac{8}{504^2(2z)^6}&\left[E_6\left(\frac{-1}{2z}\right) - E_6\left(\frac{-1}{4z}\right)\right] \\ &\times\left[E_6(z) - 33E_6(2z) + 32E_6(4z)\right].\end{aligned}$$

Suppose that $z = iu$ with $u \geq 0$. Then the right-hand side reduces to a four-dimensional infinite series. We have

$$\Delta(iu) + 24\ \Delta(2iu) + 2^{11}\ \Delta(4iu) = \frac{1}{8u^6}\sum_{\substack{n,m,r,s\geq 1\\ m,r,s \text{ odd}}}(nr)^5e^{-2\pi\left(\frac{nm}{4u}+rsu\right)}.$$

Multiply both sides by u^{k-1} and integrate for $u \in (0,\infty)$. By uniform convergence:

$$8\left(1 + 3\cdot 2^{3-k} + 2^{11-2k}\right)L(\Delta,k) = \frac{(2\pi)^k}{(k-1)!}\sum_{\substack{n,m,r,s\geq 1\\ m,r,s \text{ odd}}}(nr)^5\int_0^{\infty}u^{k-7}e^{-2\pi\left(\frac{nm}{4u}+rsu\right)}du. \tag{11}$$

Finally apply the key trick: *Use a change of variables to swap the indices of summation inside the integral in (11). If the trick is properly executed, then the right-hand side reduces to an integral over modular or quasi-modular functions.*

Theorem 1. *Recall that $F(\alpha)$ is defined in (2). Then*

$$L(\Delta,12)=-\frac{128\pi^{11}}{8241\cdot 11!}\int_0^1\left[F(\alpha)F(1-\alpha)\right]^5 \times\left(\frac{2+251\alpha+876\alpha^2+251\alpha^3+2\alpha^4}{1-\alpha}\right)\log\alpha\;\mathrm{d}\alpha. \tag{12}$$

Proof. Set $k=12$ and let $u\mapsto\frac{nu}{r}$ in (11). The formula becomes

$$\begin{aligned}\frac{8241}{2^{22}\pi^{12}}L(\Delta,12)=&\frac{1}{11!}\int_0^\infty u^5\sum_{\substack{r,m\geq1\\ r,m\text{ odd}}}\frac{1}{r}e^{-\frac{2\pi rm}{4u}}\sum_{\substack{n,s\geq1\\ s\text{ odd}}}n^{11}e^{-2\pi nsu}\mathrm{d}u\\ =&\frac{1}{11!}\int_0^\infty u^5\log\left(\prod_{\substack{m=1\\ m\text{ odd}}}^\infty\frac{1+e^{-\frac{2\pi m}{4u}}}{1-e^{-\frac{2\pi m}{4u}}}\right)\sum_{n=1}^\infty\frac{n^{11}e^{-2\pi nu}}{1-e^{-4\pi nu}}\mathrm{d}u.\end{aligned}$$

Now suppose that u and α are related by (4), and recall that $\alpha\in(1,0)$ when $u\in(0,\infty)$. We compute $\mathrm{d}u/\mathrm{d}\alpha$ using (6), and various identities from Ramanujan's notebooks imply

$$\begin{aligned}\prod_{\substack{m=1\\ m\text{ odd}}}^\infty\frac{1+e^{-\frac{2\pi m}{4u}}}{1-e^{-\frac{2\pi m}{4u}}}&=\alpha^{-1/8},\\ \sum_{n=1}^\infty\frac{n^{11}e^{-2\pi nu}}{1-e^{-4\pi nu}}&=\frac{1}{32}\alpha\left(2+251\alpha+876\alpha^2+251\alpha^3+2\alpha^4\right)\left[F(\alpha)\right]^{12}.\end{aligned}$$

Combining the formulas completes the proof. □

4. Formulas for $L(\Delta,k)$ when $k\geq 13$

It is possible to prove formulas for every value of $L(\Delta,k)$ when $k\geq 13$. Unfortunately, all of the formulas involve double integrals. A typical example is the formula for $L(\Delta,13)$ given in equation (17). It appears to be very difficult to relate these integrals to standard special functions. In fact, Zudilin's calculation of $L(\eta^2(4z)\eta^2(8z),3)$, seems to be the only instance with $k>\text{weight}(f)$, where $L(f,k)$ has been proved to equal something recognizable [11]. Boyd has given many conjectural formulas involving three-variable Mahler measures, but all of these remain open [3].

We use a variant of Zudilin's approach to study $L(\Delta,k)$, when $k\geq 13$. The calculation requires us to express a negative weight Eisenstein series in terms of an integral of modular functions. Zudilin studied a closely related Eichler integral, and obtained a formula by solving an inhomogeneous differential equation. Below we introduce a new approach for performing these calculations.

Consider equation (11). If k is even, interchange n and r using $u \mapsto \frac{nu}{r}$. If k is odd, interchange m and r using $u \mapsto \frac{mu}{r}$. Thus for k odd, the integral becomes

$$\begin{aligned} &8\frac{\left(1+3\cdot 2^{3-k}+2^{11-2k}\right)}{(2\pi)^k}L(\Delta,k) \\ &\quad = \frac{1}{(k-1)!}\int_0^\infty u^{k-7}\sum_{\substack{n,r\geq 1\\ r\text{ odd}}}\frac{n^5}{r^{k-11}}e^{-\frac{2\pi nr}{4u}}\sum_{\substack{m\geq 1\\ m\text{ odd}}}m^{k-6}\frac{e^{-2\pi mu}}{1-e^{-4\pi mu}}\mathrm{d}u. \end{aligned} \tag{13}$$

The nested sum on the right reduces to a linear combination of Eisenstein series. Unfortunately, we also have to deal with an Eisenstein series of negative weight:

$$\sum_{\substack{n,r\geq 1\\ r\text{ odd}}}\frac{n^5}{r^{k-11}}e^{-2\pi nrw}. \tag{14}$$

This main difficulty is that the sum contains negative powers of the index r. To simplify (14) consider a non-standard version of the Gamma function integral:

$$\frac{q^m}{m^s} = \frac{1}{\Gamma(s)}\int_0^q (\log q - \log v)^{s-1}\, v^m \frac{\mathrm{d}v}{v}. \tag{15}$$

After noting that (15) holds for s and m sufficiently large, we can write

$$\begin{aligned} \sum_{\substack{n,r\geq 1\\ r\text{ odd}}}\frac{n^5}{r^{k-11}}e^{-2\pi nrw} &= \sum_{\substack{n,r\geq 1\\ r\text{ odd}}} n^{k-6}\frac{e^{-2\pi nrw}}{(nr)^{k-11}} \\ &= \frac{1}{(k-12)!}\int_0^{e^{-2\pi w}} (-2\pi w - \log v)^{k-12}\sum_{n=1}^{\infty}\frac{n^{k-6}\ v^n}{1-v^{2n}}\frac{\mathrm{d}v}{v}. \end{aligned} \tag{16}$$

When k is odd, the nested sum equals a linear combination of Eisenstein series. In Theorem 2, we use (13) and (16) to obtain a formula for $L(\Delta, 13)$. Before proceeding, we note that equation (15) allows us to simplify many additional Eisenstein series of negative weight. We can use this method to simplify all of the values of $L(\Delta, k)$ when $k \geq 13$. The odd cases require formula (13), while the even cases require the analogous formula obtained from mapping $u \mapsto \frac{mu}{r}$ in (11).

Theorem 2. *Recall that $F(\alpha)$ is defined in (2). The following identity is true:*

$$\begin{aligned} &\frac{90^3\cdot 49\cdot 11\cdot 313}{\pi^{13}}L(\Delta,13) \\ &\quad = \int_0^1\int_0^\alpha \left[F(1-\beta)F(\alpha)-F(\beta)F(1-\alpha)\right]\left[F(\beta)F(\alpha)\right]^5 \\ &\qquad\qquad \times\frac{(1+\alpha)(17-32\alpha+17\alpha^2)(2+13\beta+2\beta^2)}{\alpha(1-\beta)}\mathrm{d}\beta\mathrm{d}\alpha. \end{aligned} \tag{17}$$

Proof. Combine (13) and (16) when $k=13$. We obtain

$$\frac{32865}{2^{25}\pi^{13}}L(\Delta,13)=\frac{1}{12!}\int_0^\infty u^6\sum_{\substack{m\geq 1\\ m\text{ odd}}}\frac{m^7e^{-2\pi mu}}{1-e^{-4\pi mu}}$$
$$\times\int_0^{e^{-\frac{2\pi}{4u}}}\left(-\frac{2\pi}{4u}-\log v\right)\sum_{n=1}^\infty\frac{n^7v^n}{1-v^{2n}}\frac{dv}{v}\mathrm{d}u.$$

Now suppose u and α are related by (4), and recall that $\alpha\in(1,0)$ when $u\in(0,\infty)$. We have [2, pg .133]:

$$\sum_{\substack{m\geq 1\\ m\text{ odd}}}\frac{m^7e^{-2\pi mu}}{1-e^{-4\pi mu}}=\frac{1}{64}\alpha(2-\alpha)\left(2-2\alpha+17\alpha^2\right)[F(\alpha)]^8.$$

After applying (6), the integral reduces to

$$=\frac{1}{2^{13}12!}\int_0^1\frac{(2-\alpha)\left(2-2\alpha+17\alpha^2\right)}{1-\alpha}\left[F\left(1-\alpha\right)\right]^6$$
$$\times\int_0^{q(1-\alpha)}\left(\frac{F\left(\alpha\right)}{F\left(1-\alpha\right)}+\frac{1}{\pi}\log v\right)\sum_{n=1}^\infty\frac{n^7v^n}{1-v^{2n}}\frac{\mathrm{d}v}{v}\mathrm{d}\alpha,$$

where $q(\alpha):=\exp\left(-\pi F\left(1-\alpha\right)/F\left(\alpha\right)\right)$. Next let $v=q(\beta)$, and notice that

$$\sum_{m=1}^\infty\frac{m^7v^m}{1-v^{2m}}=\frac{1}{32}\beta\left(2+13\beta+2\beta^2\right)[F(\beta)]^8,$$
$$\frac{\mathrm{d}v}{\mathrm{d}\beta}=\frac{v}{\beta(1-\beta)\left[F(\beta)\right]^2}.$$

The formula becomes

$$=\frac{1}{2^{18}12!}\int_0^1\frac{(2-\alpha)\left(2-2\alpha+17\alpha^2\right)}{1-\alpha}\left[F\left(1-\alpha\right)\right]^6$$
$$\times\int_0^{1-\alpha}\left(\frac{F\left(\alpha\right)}{F\left(1-\alpha\right)}-\frac{F\left(1-\beta\right)}{F\left(\beta\right)}\right)\frac{\left(2+13\beta+2\beta^2\right)}{1-\beta}\left[F\left(\beta\right)\right]^6\mathrm{d}\beta\mathrm{d}\alpha.$$

We conclude by letting $\alpha\mapsto 1-\alpha$. ☐

5. Conclusion

Perhaps it is regrettable that our formulas for $L(\Delta,k)$ are complicated. While the formulas demonstrate that $L(\Delta,k)$ belongs to the ring of periods for $k\geq 12$, it would be nice if they could be simplified to a recognizable form. We would find it particularly exciting if $L(\Delta,12)$ could be related to special values of generalized hypergeometric functions. We are also willing to speculate that there might be a relation between $L(\Delta,12)/\pi^{12}$ and the Mahler measure of some 12-variable polynomial.

Acknowledgements. The author thanks Wadim Zudilin for his frequent encouragement and many useful suggestions.

References

[1] A. Beilinson, Higher regulators of modular curves, in: *Applications of Algebraic K theory to Algebraic Geometry and Number Theory.* Contemp. Math. **55**, Amer. Math. Soc., Providence (1986),1–34.

[2] B. C. Berndt, *Ramanujan's Notebooks, Part III* (Springer-Verlag, New York, 1991).

[3] D. Boyd, Mahler's measure and special values of L-functions of elliptic curves at $s = 3$, Preprint (2006). `http://www.math.ubc.ca/~boyd/sfu06.ed.pdf`

[4] M. Kontsevich and D. Zagier, Periods, in: *Mathematics unlimited — 2001 and beyond* (Springer, Berlin, 2001), 771–808.

[5] K. Ono, The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and q-series, *American Mathematical Society, Providence, RI,* 2004.

[6] S. Ramanujan, On certain Arithmetical Functions, [Trans. Camb. Phil. Soc. **2** (1916), no. 9, 159–184]. *Collected papers of Srinivasa Ramanujan, 23-29, AMS Chelsea Publ., Providence, RI, 2000.*

[7] R. A. Rankin *Modular Forms and Functions*, Cambridge University Press, Cambridge, 1977.

[8] M. Rogers and W. Zudilin, From L-series of elliptic curves to Mahler measures, *Compositio Math.* **148** (2012), no. 2, 385-414.

[9] M. Rogers and W. Zudilin, On the Mahler measure of $1 + X + 1/X + Y + 1/Y$, Preprint arXiv:1102.1153 [math.NT] (February 2011), 18 pages.

[10] D. Zagier Introduction to modular forms. in: *From Number Theory to Physics*, M. Waldschmidt et al, Springer-Verlag, Heidelberg (1992) 238–291.

[11] W. Zudilin, Period(d)ness of L-values, *Proceedings of the International Conference in memory of Alf van der Poorten (to appear)*, J.M. Borwein et al. (eds.), 14 pages.

Department of Mathematics and Statistics, Université de Montréal, CP 6128 succ. Centre-ville, Montréal Québec H3C 3J7, Canada

E-mail address: `mathewrogers@gmail.com`